



# Chlorpyrifos Factsheet

---

## What is chlorpyrifos?

Chlorpyrifos is an organophosphate insecticide used in agriculture primarily on cotton, oranges, corn, and almonds, among many other crops. Also known as Lorsban (agricultural use products), among other trade names, and manufactured predominantly by Dow AgroSciences, chlorpyrifos is one of the most widely used insecticides in the U.S. Nationwide in 2001 (prior to the cancellation of residential uses) US EPA estimated that 11–16 million pounds of the insecticide were used, second only to malathion for insecticide use.<sup>1</sup>

Chlorpyrifos was widely used in homes as a residential insecticide (under the trade name Dorsban) through 2005 when its phaseout for those uses was completed—the result of a U.S. Environmental Protection Agency regulatory decision based primarily on its hazards for children.<sup>2</sup> Although some agricultural uses were altered by this phaseout agreement—to reduce residue levels in harvested produce, agricultural use remains high. In California in 2004, 2.3 million pounds of chlorpyrifos were reported sold.<sup>3</sup>

## Exposure to chlorpyrifos

People can be exposed to chlorpyrifos by direct dermal contact, contact with treated surfaces, ingestion of chlorpyrifos contaminated dust, and by breathing air inside or outside treated buildings or near fields where it was applied. Exposures also result from eating food contaminated with chlorpyrifos residues.<sup>4,5</sup> Contact with flea collars containing chlorpyrifos can also lead to exposure.

Chlorpyrifos is found in particularly high levels in children. Data from the Center for Disease Control and Prevention (CDC) showed that 93% of U.S. residents sampled between 1999 and 2000 had chlorpyrifos in their bodies. Children (6–11 years) showed levels almost twice as high as those of adults. Chronic exposure levels were 4.6 times the “acceptable” level for children and 3.0 times the “acceptable” level for youth (12–19 years).<sup>6</sup>

## Health Effects

Organophosphate pesticides’ most well-known and documented effects are their inhibition of an enzyme (acetyl cholinesterase), necessary for proper transmission of nerve impulses.<sup>7</sup> Symptoms of low-dose exposure may include headaches, agitation, inability to concentrate, weakness, tiredness, nausea,

diarrhea and blurred vision. At higher doses, abdominal cramps, vomiting, sweating, tearing, muscular tremors, pinpoint pupils, low blood pressure, slow heartbeat and difficulty breathing may be observed.<sup>8</sup>

The Association of Occupational and Environmental Clinics (AOEC) lists all organophosphates generally and chlorpyrifos specifically as capable of causing asthma in previously unaffected individuals.<sup>9</sup> Exposure can also exacerbate asthmatic symptoms in individuals who already have the disease.

In addition to acute symptoms, many recent studies indicate that low-level exposure to chlorpyrifos interferes with the development of the mammalian nervous system.<sup>10</sup> Neural cell replication and differentiation are both affected, with a reduction in the number of neural connections observed in exposed rats.<sup>11</sup>

Human epidemiological studies on pregnant women exposed to chlorpyrifos through involuntary home pesticide use demonstrate a link between *in utero* exposure to chlorpyrifos and low birth weights and reduced head circumference of newborns. These effects were most significant for individuals whose genetic makeup is such that they produce low levels of PON1, the enzyme responsible for detoxifying chlorpyrifos in the body.<sup>12</sup> Chlorpyrifos is also a suspected endocrine disrupting compound; moderate doses have been shown to alter hormone levels in animal studies.<sup>13</sup>

In addition to heightened vulnerability to chlorpyrifos because of their developing nervous systems, children are likely to be exposed to higher levels of chlorpyrifos than adults for several reasons. Children eat, breathe, and drink more per pound of body weight than adults. Children also play on the floor and in the grass where pesticide residues collect; and they exhibit hand-to-mouth behaviors that increase their potential for exposure.

While the U.S. EPA lists chlorpyrifos as providing evidence of no carcinogenicity, recent studies suggest possible links to both lung<sup>14</sup> and prostate<sup>15</sup> cancer.

## Chlorpyrifos is a significant air pollutant

Chlorpyrifos is a semi-volatile chemical that readily leaves in gas form from leaf and soil surfaces to

become airborne, especially when outdoor temperatures are high. A study done in July and August 2004 and 2005 in Lindsay, California, demonstrated the presence of chlorpyrifos in the air near homes in this agricultural community. Over 100 air samples were collected near homes and three-quarters of the samples had detectable levels. Eleven percent of the samples were above the levels determined to be “acceptable” for a 24-hour exposure by children. The highest concentration observed was nearly eight times the allowable level.<sup>16</sup> Air monitoring studies conducted by the California Air Resources Board also indicated the potential for high exposures in areas near application sites.<sup>17</sup>

## Ecological Effects

Chlorpyrifos is moderately to very highly toxic to birds, honeybees, fish, and aquatic organisms. According to the EPA, just a single application of chlorpyrifos poses a risk to small mammals as well.<sup>18</sup> It is a persistent chemical that accumulates in the tissue of aquatic organisms and has a half-life in soil ranging from 60 days to one year depending on climate and soil type.<sup>19</sup>

*Revised October 2006*

- 
- 1 U.S. Environmental Protection Agency. Pesticide Industry Sales and Usage: 2000 and 2001 Market Estimates. Available at <http://www.epa.gov/oppbead1/pestsales/index.htm>.
  - 2 Chlorpyrifos Revised Risk Assessment and Agreement with Registrants, U.S. EPA, June 2000, <http://www.epa.gov/pesticides/op/chlorpyrifos/agreement.pdf>.
  - 3 CA Pesticide Sales Data for 2004, California Department of Pesticide Regulation, Sacramento, CA, <http://www.cdpr.ca.gov/docs/mlsassess/nopds/old.htm>. The 2004 data are the most recent available at the time this report was written.
  - 4 Agency for Toxic Substances and Disease Registry. ToxFAQs(tm) for Chlorpyrifos. September 1997. Available at <http://www.atsdr.cdc.gov/tfacts84.html>.
  - 5 Fenske, RA, L. Chensheng, D. Barr, and L. Needham. Children’s exposure to Chlorpyrifos and Parathion in an agricultural community in central Washington state. *Environmental Health Perspectives* 2002; 110(5):549-553.
  - 6 Schafer, K.S., M. Reeves, S. Spitzer and S.E. Kegley. Chemical Trespass: Pesticides in our bodies and corporate accountability. Pesticide Action Network, San Francisco, CA. 2004. <http://www.panna.org/campaigns/docs/Trespass/chemicalTrespass2004.dv.html>.
  - 7 Toxicology Information Briefs: Cholinesterase Inhibition, Extension Toxicology Network, Cooperative Extension Offices of Cornell University, Oregon State University, the University of Idaho, and the University of California at Davis and the Institute for Environmental Toxicology, Michigan State University. <http://extoxnet.orst.edu/tibs/cholines.htm>.
  - 8 Reigert, JR and Roberts, JR, Recognition and Management of Pesticide Poisoning, 5th ed., US EPA, 1999, <http://www.epa.gov/pesticides/safety/healthcare/handbook/contents.htm>.
  - 9 AOEC Exposure Codes, George Washington University, Occupational Medicine Group for the Association of Occupational and Environmental Clinics, <http://www.aoec.org/aoeccode.htm>.
  - 10 Slotkin, TA, Levin, ED, Seidler, FJ, Comparative Developmental Neurotoxicity of Organophosphate Insecticides: Effects on Brain Development Are Separable from Systemic Toxicity, *Envi. Health Persp.* 2006, 114(5): 746-51 and references cited therein.
  - 11 a) Aldridge JE, Seidler FJ, Meyer A, Thillai I, Slotkin TA, Serotonergic Systems Targeted by Developmental Exposure to Chlorpyrifos: Effects during Different Critical Periods. *Envi. Health Persp.* 2003, 111(14): 1736-43. b) Jameson, RR, Seidler, FJ, Qiao, D, Slotkin, TA, Chlorpyrifos Affects Phenotypic Outcomes in a Model of Mammalian Neurodevelopment: Critical Stages Targeting Differentiation in PC12 Cells, *Envi. Health Persp.* 2006, 114(5): 667-72. c) Aldridge, JE, Meyer, A, Seidler, FJ, Slotkin, TA, Alterations in Central Nervous System Serotonergic and Dopaminergic Synaptic Activity in Adulthood after Prenatal or Neonatal Chlorpyrifos Exposure, *Envi. Health Persp.* 2005, 113(8): 1027-31.
  - 12 a) Whyatt RM and Barr, DB, Measurement of organophosphate metabolites in postpartum meconium as a potential biomarker of prenatal exposure: A validation study, *Envi Health Persp.* 2001, 109(4): 417-20. b) Whyatt RM, Rauh V, Barr DB, et al. Prenatal Insecticide Exposures and Birth Weight and Length among an Urban Minority Cohort, *Envi. Health Persp.* 2004, 112(10):1125-32. c) Berkowitz GS, Wetmur JG, Birman-Deych E, et al., In Utero Pesticide Exposure, Maternal Paraoxonase Activity, and Head Circumference, *Envi. Health Persp.* 2004, 112(3):388-91. (d) Furlong, CE, Holland N, Richtera RJ, Bradman A, Ho A and Eskenazi B. PON1 status of farmworker mothers and children as a predictor of organophosphate sensitivity. *Pharmacogenetics and Genomics* 2006, 16:183-190.
  - 13 a) Rawlings, NC, Cook, SJ, and Waldbillig, D, Effects of the pesticides carbofuran, chlorpyrifos, dimethoate, lindane, triallate, trifluralin, 2,4-D, and pentachlorophenol on the metabolic endocrine and reproductive endocrine system in ewes, *J Tox. Envi. Health*, 1998, 54: 21-36. b) Keith, LH, *Environmental Endocrine Disruptors: A Handbook of Property Data*, Wiley Interscience (New York, 1997).
  - 14 Lee, WJ, A. Blair, JA Hoppin, JH Lubin, JA Rusiecki, DP Sandler, M Dosemeci, MC Alavanja. Cancer incidence among pesticide applicators exposed to chlorpyrifos in the Agricultural Health Study. *J Natl Cancer Inst.* 1:96(23):1781-9. 2004.
  - 15 Alavanja, MCR, C Samanic, M Dosemeci, J Lubin, R Tarone, CF Lynch, C Knott, K Thomas, JA Hoppin, J Barker, J Coble, DP Sandler and A Blair. Cancer: possible association with increase risk of prostate cancer. Use of Agricultural Pesticides and Prostate Cancer Risk in the Agricultural Health Study Cohort. *Am J of Epidem* 157:800-814. 2003.
  - 16 Mills K and S Kegley. Air Monitoring for Chlorpyrifos in Lindsay, California: June-July 2004 and July-August 2005. Pesticide Action Network, San Francisco, CA. April 2006.
  - 17 Report for the Application and Ambient Air Monitoring of Chlorpyrifos (and the Oxon Analogue) in Tulare County during Spring/Summer 1996, California Air Resources Board, Test Report #C96-040 and # C96-041, April 7, 1998, <http://www.cdpr.ca.gov/docs/emppm/pubs/tac/chlrpfs.htm>.
  - 18 See reference #2
  - 19 Extension Toxicology Network. “Chlorpyrifos.” June 1996. Available at <http://extoxnet.orst.edu/pips/chlorpyrifos.htm>.
-